To patients prescribed with INAVIR and their carers.

Take a deep breath and relax. It is not necessary to inhale excessively, nor to hold your breath for a long time.





Open the package just before use.






INAVIR is an inhaled (oral inhalation) medicine for the treatment of influenza.

INAVIR is an inhalation medicine with antiviral effects to prevent the influenza virus from spreading inside your body. (The package contains powdered medication)







Benefits of inhaled medication The influenza virus multiplies in the "throat" and "windpipe" When inhaled, INAVIR is delivered directly to the throat and windpipe where the virus Inhale multiplies, and prevents the virus from multiplying. INAVIR is delivered directly to the "throat" and "windpipe"

**Carers should supervise children prescribed with INAVIR, as children may find the Medication Tray difficult to slide.

Video instructions on how to inhale and information about influenza are available on "Influ News."



http://www.influ-news.info




When using INAVIR, you

"slide and inhale".

"Slide and inhale" Medicine Tray 1 and 2.







Inhale

Before inhaling INAVIR... You can practice inhaling using the inhaler.

You can practice inhaling by using the inhaler but without sliding the Medicine Trav.



Stay relaxed when you inhale INAVIR.

Slide 0



Without sliding the Medicine Tray, tap the container gently onto a table, "tap, tap, tap". This is to drop the medicine to the bottom of the inhaler.

Inhale



Without removing the labels, slide Medicine Tray 1 in the direction of the arrow all the way to the end.

Put the mouthpiece into your mouth and take a deep breath, then hold your breath for 2 - 3 seconds. Remove the inhaler from vour mouth and exhale slowly without blowing into the mouthpiece.

Slide 2



Next, slide Medicine Tray in the direction of the arrow all the way to the end.

Inhale



Put the mouthpiece into your mouth and take a deep breath, then hold your breath for 2 - 3 seconds. Remove the inhaler from vour mouth and exhale slowly without blowing into the mouthpiece.

Return to original position



Slide Medicine Tray 1 back to its original position.

F

To ensure that the medicine is completely inhaled. repeat procedures A-C.

Tips ģ inhaling INAVIR

Instructions

on how

6

use

INAVIR



Do not tap the inhaler after sliding the Medicine Trav.





Do not cover the air hole.



Put the mouthpiece fully into your mouth.



Breathe out briefly before putting the mouthpiece into your mouth. This will make it easier to inhale.



Inhale in an upright position, not looking down.



Hold your breath for 2 - 3 seconds after inhaling, then breathe out slowly.

* Some people might find the medicine bitter.



and dose of INAVIR

turn over for usage

